

Outline

- Product Quality Research Institute (PQRI) Stability Shelf Life Working Group
- Shelf Life Estimation
- Simulation Results
- Example using real-life data
- Future/Continued Research

Product Quality Research Institute (PQRI) Stability Shelf Life Working Group

- Objectives
 - Investigate statistical methods for estimating shelf life which allow the sponsor to define and manage risk
 - Assess alternative methods for estimating shelf life
 - Enhance safety/efficacy of pharmaceutical products through accurate estimation of shelf life
- Research efforts include developing statistical methodology to directly estimate shelf life of pharmaceutical products

Shelf Life Estimation

- ICH Guidelines
 - Q1E states the purpose of a stability study is to establish “a retest period or shelf life and label storage instructions applicable to all future batches manufactured and packaged under similar circumstances”
 - Test for equal slopes/intercepts among batches using $\alpha = 0.25$
 - No evidence of batch-to-batch variability: labeled shelf-life is the time the 95% one-sided lower (upper) confidence bound for mean degradation curve/stability limiting characteristic intersects lower (upper) specification limit
 - Evidence of batch-to-batch variability: shelf-life for each individual batch is computed and minimum of all shelf-lives is labeled shelf-life
- Problems with ICH Guidelines
 - Shao & Chow (1994) conclude the minimum approach “lacks statistical justification”
 - Estimate obtained from pooling batches applies only to batches used in analysis; inference should be made to future batches


Random Batch Analysis

- Proposed statistical analysis takes into account batch-to-batch variation via random batch effects
- By appropriately accounting for batch-to-batch variation, the question of batch “poolability” is eliminated
- Including batch-to-batch variation allows for more straightforward estimation and interpretation of shelf life
- Inference can be made to future batches


Proposed Methodology

- Provides a consistent, flexible methodology for directly estimating shelf life
 - Consistent with how acceptance criteria is defined
 - Can be implemented using overall mean response or percentile of response distribution
- Use mixed model (random batch effects) on stability limiting response
 - Estimated shelf life is then “applicable to all future batches” (ICH Q1E)
 - Involves lower interval estimate on calibrated point

Proposed Shelf Life Estimation Procedure



Proposed Shelf Life Estimation Procedure



- Estimated shelf life is storage time corresponding to the point where predicted mean (or quantile) response intersects specification limit or acceptance criteria
- Lower interval estimate is constructed around calibrated point to determine labeled shelf life
 - Similar to Shao & Chow’s (1994) $1-\alpha$ lower confidence bound for ε^{th} quantile of true shelf life
- As added information on the quality of the labeled shelf life estimate, 2-sided interval estimate (e.g. CI, PI, TI) is obtained about labeled shelf life
 - 2-sided interval estimate is a diagnostic tool
 - Analogous to Chow’s (2007) *safety margin* which provides useful information regarding drug safety beyond labeled shelf life
 - Similar to Chow & Shao’s (1991) tolerance correction to estimate the 95% lower bound for individual shelf lives

Interval Estimates on Calibrated Point

- 3 methods to obtain interval estimate about calibrated point:
 - Using distribution of x_0
 - Using distribution of estimated parameter values ($\hat{\beta}$)
 - Reflection method
- Method 2 produces less conservative estimates (usually \geq other methods in linear case)

Simulation Example

- Data simulated for 36 months using 3 and 6 batches
- Acceptance criteria is 95-105% of label claim
- Assay follows simple linear response decay
- Model: $y_{ij} = b_0 + batch_{i0} + (b_1 + batch_{i1})x_{ij} + \varepsilon_{ij}$
 - $batch_{i0}$ = random batch effect on intercept
 - $batch_{i1}$ = random batch effect on slope
 - ε_{ij} = error associated with response for i^{th} batch, j^{th} month
 - $\varepsilon_{ij} \sim N(0, \sigma_{\varepsilon}^2)$ $batch_{i0} \sim N(0, \sigma_b^2)$ $batch_{i1} \sim N(0, \sigma_{b,m}^2)$
- NLMIXED was used to analyze mean response and is compared with ICH approach

Simulation Results (3, 6 batches), $\alpha = 0.05$

- Percentage of time true shelf life is captured ($labeled \leq true$)
 - ICH method: **98%, 99%**
 - Mixed Model method: **98%, 95%**
- Average difference between true and estimated shelf life
 - ICH: **4.7 months, 5.7 months**
 - Mixed Model: **4.6 months, 2.4 months**

Simulation Results – 6 batches True Shelf Life: 33.3 months

Method	Sim. #	Average	Avg. diff	Under %	Over %	Under diff	Over diff	Min	Max
ICH	1	27.5	-5.8	0.997	0.003	-5.9	0.9	20.5	35.4
ICH	2	27.6	-5.7	0.998	0.002	-5.7	0.2	20.3	33.6
ICH	3	27.6	-5.7	0.999	0.001	-5.8	0.3	21.0	33.7
ICH	4	27.7	-5.7	0.999	0.001	-5.7	0.3	22.0	33.6
ICH	5	27.6	-5.7	0.998	0.002	-5.8	0.5	19.2	34.2
LMM_B_hat	1	30.9	-2.4	0.951	0.049	-2.6	0.6	26.6	36.2
LMM_B_hat	2	31.0	-2.3	0.948	0.052	-2.5	0.6	26.2	35.4
LMM_B_hat	3	31.0	-2.3	0.945	0.055	-2.5	0.7	25.8	36.2
LMM_B_hat	4	31.1	-2.3	0.941	0.059	-2.5	0.7	27.0	36.3
LMM_B_hat	5	31.1	-2.3	0.945	0.055	-2.5	0.7	27.0	35.9
LMM_Reflect/x_0	1	30.8	-2.6	0.961	0.039	-2.7	0.6	26.9	36.0
LMM_Reflect/x_0	2	30.9	-2.4	0.951	0.049	-2.6	0.5	26.5	35.2
LMM_Reflect/x_0	3	30.9	-2.4	0.949	0.051	-2.6	0.6	25.7	36.1
LMM_Reflect/x_0	4	30.9	-2.4	0.951	0.049	-2.6	0.6	26.9	36.1
LMM_Reflect/x_0	5	30.9	-2.4	0.952	0.048	-2.6	0.7	26.9	35.8

Adding more batches

- ICH method:
 - increases bias (farther away from true shelf life)
 - overestimation rate approaches 0
- Mixed Model method:
 - decreases bias (closer to true shelf life)
 - overestimation rate approaches α
- On average Mixed Model method produces longer, more accurate shelf life


Simulation Results, True Shelf Life: 33.3 months

- 3 batches: ICH and Mixed Model methods produce on average equal estimated shelf lives, but estimate is not good (\ll true shelf life)
- 6 batches: ICH method: **27.6 months** Mixed Model method: **30.9 months**
- 9 batches: ICH method: **26.9 months** Mixed Model method: **31.4 months**
- 12 batches: ICH method: **26.5 months** Mixed Model method: **31.7 months**
- Do we want an estimator whose bias increases as $n \rightarrow \infty$ (ICH) or whose bias $\rightarrow 0$ as $n \rightarrow \infty$ (Mixed Model) ???


Example using 6 batches of real-life data

- Data:
 - Blinded
 - 24-month
 - assay response (% label claim)
 - specification limits 90-110%
- Shelf life estimate using ICH guidelines: **23.8 months**
- Shelf life estimate using Mixed Model: **28 months**

Real-life data (using ICH)



Real-life data (using Mixed Model)



Future/Continued Research

- Develop theory/methodology for quantile regression with random batch effects to estimate shelf life
 - Model a quantile of response distribution instead of mean
- Determine robustness of proposed method to estimate shelf life using a limited number of months of real-life data
- Determine the optimal interval to construct around labeled shelf life
- Determine sampling distribution of estimates using ICH and proposed methodology

Acknowledgements

This research is funded through the PQRI Stability Shelf Life Working Group. PQRI is the Product Quality Research Institute, Arlington, Virginia